




BounceBack[®] Foliar 4-0-0

FOLIAR-APPLIED LIQUID NUTRIENT

N 4%

Ca 8%



4-0-0

GUARANTEED ANALYSIS

Total Nitrogen (N) 4.0%
4.0% Nitrate Nitrogen
Calcium (Ca) 8.0%

Derived from: calcium gluconate and calcium nitrate.

2.5 US Gal / 9.5 Liters
12.0 lbs per Gallon @ 68° F
1.4 kgs per Liter @ 20° C
Net Weight 30.0 lbs / 13.6 kgs

DO NOT STORE BELOW 40°F (5°C)

FBSciences Inc.

153 N. Main St, Suite 100 | Collierville, TN 38017 USA

USA092722 | F1837

3102410

BounceBack[®] Foliar 4-0-0

DIRECTIONS FOR USE

DO NOT spray in the heat of the day or when the plant is under moisture stress. DO NOT spray to the point of runoff. Use as fine of a spray mist as possible. Allow 3-4 hours after application before sprinkler irrigating to avoid washing the product off.

Recommended mixing sequence: water, adjuvants, pesticides, FBSciences nutrient products, other fertilizers, balance of water while agitating. When mixing with high phosphate fertilizers, add a citric acid buffer until the pH is 4.5 to 5.0 to improve compatibility and uptake. Ensure agitation is available when mixing with sulfate fertilizers. A standard jar test is recommended before tank mixing.

RATES & TIMING FOR ALL CROPS

Apply **1-6 quarts/acre** anytime during the growing season; repeat as needed. The higher rates should be used for orchards and vineyards when using spray volume greater than 50 gallons per acre.

INSTRUCCIONES DE USO

NO rociar en el calor del día o cuando la planta esté bajo estrés por humedad. NO rociar hasta el punto de escorrentía. Usar una niebla de rociado lo más fina posible. Esperar de 3 a 4 horas después de la aplicación antes del riego por aspersión para evitar lavar el producto.

Secuencia de mezcla recomendada: agua, adyuvantes, pesticidas, productos nutritivos de FBSciences, otros fertilizantes, balance de agua mientras se agita. Cuando se mezcle con fertilizantes de alto contenido de fósforo, agregar un regulador de ácido cítrico hasta que el pH esté de 4.5 a 5.0 para mejorar la compatibilidad y la absorción. Asegúrese de que la agitación esté disponible al mezclar con fertilizantes de sulfato. Se recomienda una prueba de jarra estándar antes de mezclar el tanque.

TASAS Y MOMENTO PARA TODOS LOS CULTIVOS

Aplicar de **1-6 cuartos de galón/acre** en cualquier momento durante la temporada de crecimiento; Repetir según sea necesario. Las dosis más altas deben usarse para huertos y viñedos cuando se usa un volumen de rociado mayor a 50 galones por acre.

WARNING

READ ENTIRE LABEL BEFORE USE | KEEP OUT OF REACH OF CHILDREN



HAZARD STATEMENTS Harmful if swallowed. Causes serious eye irritation.

AVISO PELIGRO Dañino si se ingiere. Causa irritación ocular grave.

PRECAUTIONARY STATEMENTS Wear protective gloves/clothing and eye/face protection. Wash skin thoroughly after handling. Do not eat, drink or smoke when using this product. **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If eye irritation persists, get medical attention. **IF SWALLOWED:** Rinse mouth. Call a POISON CENTER or doctor/physician if you feel unwell. Dispose of contents/container in accordance with local/national regulations.

WARRANTY AND CONDITIONS OF SALE Manufacturer warrants that this material conforms to the chemical description hereon and is reasonably fit for use as directed. No other warranty is made, either expressed or implied. BEFORE OPENING, see other conditions and limitations that apply which are available at: <http://www.FBSciences.com/ProductConditionsOfSale/>. See patent and other patents pending: <http://www.fbsciencespatents.com>. Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.aapfco.org/metals.html>

USA092722 | 3152410