



Phosron[®] Foliar

FOLIAR-APPLIED LIQUID NUTRIENT

N 7%

P₂O₅ 21%

K₂O 3%

Mo 0.2%

7-21-3

GUARANTEED ANALYSIS

Total Nitrogen (N)	7.0%
6.5% Ammoniacal Nitrogen	
0.5% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	21.0%
Soluble Potash (K ₂ O)	3.0%
Molybdenum (Mo)	0.2%

Derived from: ammonium hydroxide, urea, phosphoric acid, potassium hydroxide, and sodium molybdate.

2.5 US Gal / 9.5 Liters

10.7 lbs per Gallon @ 68° F

1.3 kgs per Liter @ 20° C

Net Weight 26.7 lbs / 12.1 kgs



DO NOT STORE BELOW 40°F (5°C)

FBSciences Inc.

153 N. Main St, Suite 100 | Collierville, TN 38017 USA

USA080322 | F1837

3102476



Phosron® Foliar

DIRECTIONS FOR USE

DO NOT spray in the heat of the day or when the plant is under moisture stress. DO NOT spray to the point of runoff. Use as fine of a spray mist as possible. Allow 3-4 hours after application before sprinkler irrigating to avoid washing the product off.

Recommended mixing sequence: water, adjuvants, pesticides, FBSciences nutrient products, other fertilizers, balance of water while agitating. When mixing with calcium or micronutrient fertilizers, add a citric acid buffer until the pH is 4.5 to 5.0 to improve compatibility and uptake. Ensure agitation is available when mixing with calcium fertilizers. Please consult your local FBS Representative for any questions. A standard jar test is recommended before tank mixing.

RATES & TIMING FOR ALL CROPS

Apply **2-5 quarts/acre** at any growth stage. When using spray volumes of 50 gallons per acre or greater, use the higher labeled spray rates. Repeat every two to four weeks during periods of high phosphorus demand by the crop.

WARNING: The application of fertilizer material containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant animals.

WARNING

READ ENTIRE LABEL BEFORE USE | KEEP OUT OF REACH OF CHILDREN



HAZARD STATEMENTS Causes skin and eye irritation.

AVISO PELIGRO Provoca irritación cutánea y ocular.

PRECAUTIONARY STATEMENTS Wash hands thoroughly after handling. Wear protective gloves. **IF ON SKIN:** Wash with plenty of soap and water. If skin irritation occurs, get medical advice/attention. Take off contaminated clothing and wash before reuse. **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If eye irritation persists, get medical advice/attention.

WARRANTY AND CONDITIONS OF SALE Manufacturer warrants that this material conforms to the chemical description hereon and is reasonably fit for use as directed. No other warranty is made, either expressed or implied. BEFORE OPENING, see other conditions and limitations that apply which are available at: <http://www.FBSciences.com/ProductConditionsOfSale/>. See patent and other patents pending: <http://www.fbsciencespatents.com>. Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.aapfco.org/metals.html>

INSTRUCCIONES DE USO

NO rociar en el calor del día o cuando la planta esté bajo estrés hídrico. NO rociar hasta el punto de escurrimiento. Usar una neblina de rociado tan fina como sea posible. Deje pasar 3-4 horas después de la aplicación antes de regar con aspersión para evitar lavar el producto.

Secuencia de mezcla recomendada: agua, adyuvantes, pesticidas, productos nutritivos de FBSciences, otros fertilizantes, balance de agua mientras se agita. Al mezclar con fertilizantes de calcio o micronutrientes, agregar un buffer de ácido cítrico hasta que el pH esté de 4.5 a 5.0 para mejorar la compatibilidad y la absorción. Asegurarse de que la agitación este disponible cuando se mezcla con fertilizantes de calcio. Consulte a su representante local de FBS si tiene alguna pregunta. Se recomienda realizar una prueba de jarra estándar antes de mezclar en el tanque.

TASAS Y MOMENTO PARA TODOS LOS CULTIVOS

Aplicar de **2 a 5 cuartos de galón/acre** en cualquier etapa de crecimiento. Cuando al utilizar volúmenes de rociado de 50 galones por acre o más, utilizar las tasas de rociado más altas en la etiqueta. Repetir cada dos a cuatro semanas durante los períodos de alta demanda de fósforo por parte del cultivo.